

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : A61K 31/425, C07D 277/66, 263/56, A61K 31/42		A1	(11) International Publication Number: WO 95/06469 (43) International Publication Date: 9 March 1995 (09.03.95)
(21) International Application Number: PCT/GB94/01883 (22) International Filing Date: 30 August 1994 (30.08.94) (30) Priority Data: 9317949.7 28 August 1993 (28.08.93) GB (71) Applicant (for all designated States except US): CANCER RESEARCH CAMPAIGN TECHNOLOGY LIMITED [GB/GB]; Cambridge House, 6-10 Cambridge Terrace, Regent's Park, London NW1 4JL (GB). (72) Inventors; and (75) Inventors/Applicants (for US only): STEVENS, Malcolm, Francis, Graham [GB/GB]; Shepshed Fields Farm, Remp- stone Lane, Belton, Leicestershire LE12 9XA (GB). McCALL, Carol, Jane [GB/GB]; Chestnut Cottage, 22 High Street, Pulloxhill, Bedfordshire MK45 5HA (GB). LELIEVELD, Petrus [NL/NL]; Johan Frisoplantsoen 14, NL-2751 XR Moerkapelle (NL). (74) Agent: HLN. & W.S. SKERRETT; Charles House, 148/9 Great Charles Street, Birmingham B3 3HT (GB).			(81) Designated States: AU, CA, JP, US, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE). Published <i>With international search report.</i>
(54) Title: BENZAZOLE COMPOUNDS FOR USE IN THERAPY			
(57) Abstract There are disclosed herein benzazole compounds, exemplified by 2-(4-aminophenyl)benzothiazole and analogues or salts thereof, which exhibit very significant selective cytotoxic activity in respect of tumour cells, especially breast cancer cells, and which provide potentially useful chemotherapeutic agents for treatment of breast cancer.			

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgyzstan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Latvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

Benzazole compounds for use in therapy

Field of the Invention

5 The present invention relates to certain biologically active benzazole compounds, particularly benzothiazole and benzoxazole compounds, which are able selectively to inhibit proliferation of certain mammalian tumour cells, particularly breast cancer cells, and which are accordingly
10 of interest for use as active chemotherapeutic agents in antitumour therapy, especially in connection with the treatment of breast cancer.


Summary of the Invention

15

 In one aspect this invention provides, for use in therapy, 2-arylbenzazole compounds that are especially active in inhibiting proliferation of certain breast cancer tumour cells, said compounds being exemplified by 2-(4-aminophenyl)benzothiazole and close analogues or acid
20 addition salts thereof.

 More specifically, the 2-arylbenzazole compounds of the present invention are generally compounds having the
25 structural formula I,

30



I

35 characterised in that

X is S or O;

R¹ and R³ are each independently hydrogen, alkyl, hydroxyl, alkoxy or aralkoxy

Z is aryl;

subject to the proviso that alkyl groups when present as such in the compound or as a moiety in other groups such as alkoxy are each composed of less than 6 carbon atoms;

5


or are compounds that are pharmaceutically acceptable salts of compounds represented by the structural formula specified above.

10 In addition to phenyl or substituted phenyl aromatic groups, the term aryl as used herein may also include arylalkyl (aralkyl) or aromatic heterocyclic groups, particularly nitrogen-containing heterocyclic groups such as pyridyl. Arylalkyl groups can include groups such as
15 benzyl or cinnamyl for example, and in general the aryl groups will contain at least one nitrogen atom or nitrogen containing substituent, especially an amine or a group readily convertible into an amine.

20 The invention also includes compounds that in use can act as precursors or prodrugs and readily break down or be converted, for example by metabolic processes, in the animal body or other biological systems so as to form the particular biologically active compounds hereinbefore
25 specified, especially 2-arylbenzazole compounds in which the aryl group has an amino group substituent. The claims appended hereto should therefore be construed accordingly in determining the scope of the invention.

30 In at least most preferred embodiments of the invention wherein the benzazole compound is as specified above, the group Z may be represented by the structural formula

35



II

where $n = 0$ or 1 ;


R^2 is hydrogen, NH_2 , NO_2 ,
 N_3 , halogen or an alkyl or
substituted alkyl oxysulphonyl
group;

R^4 is hydrogen, NO_2 , N_3 , pyrrolidino,
piperidino, morpholino or
 NR^5R^6 where

R^5 and R^6 each represent
hydrogen or alkyl;

with the further proviso that R^2 and R^4 are not both
hydrogen. Also, at least one of R^2 and R^4 will generally
be an amino or a substituted amino group, or some other
nitrogen-containing group convertible into an amino group.
Furthermore, usually, but not necessarily, R^4 will be para
or in the 4-position of the phenyl group.

Particularly preferred embodiments include at least
some compounds in which Z can be represented by the
structural formula



III

where R^2 is as previously defined. If R^2 is halogen, often
it will preferably be a 2-Cl group.

In some compounds within the scope of the invention,
it is also possible for the group $-(CH_2)_n-$ in the above
structure II to be replaced by an alkenylic group such as
 $-CH=CH-$

Preferred compounds of the invention in accordance
with formula I wherein R^3 is hydrogen include compounds in

which R^1 is alkoxy or benzyloxy, and X is preferably sulphur. More generally, preferred compounds of the invention in accordance with the structural formula I may be further characterised by at least one of the following 5 features:

- (a) at least some alkyl groups when present as such or as a moiety in other groups such as alkoxy are methyl or ethyl;
- 10 (b) halo substituents, when present, are selected from fluorine and chlorine;
- (c) R^2 is hydrogen or 2-halogen, and R^4 is amino.

Compounds in accordance with the invention which 15 conform to formula I wherein Z has the structural formula II with $n=0$, and which are of particular interest, include those compounds where the combination of substituents R^1 , R^2 , R^3 , R^4 and X is selected from the following combinations:

20

	<u>R^1</u>	<u>R^3</u>	<u>X</u>	<u>R^2</u>	<u>R^4</u>
	H	H	S	H	4-NH ₂
	H	H	S	3-NH ₂	4-H
25	H	H	S	2-NH ₂	4-H
	H	H	S	H	4-NMe ₂
	H	H	S	H	4-NEt ₂
	H	H	S	H	4-Pyrrolidino
30	H	H	S	H	4-Piperidino
	H	H	S	H	4-Morpholino
	H	H	S	H	4-NO ₂
	H	H	S	3-NO ₂	4-H
	H	H	S	2-NO ₂	4-H
35	H	H	S	H	4-N ₃
	H	H	S	3-N ₃	4-H
	H	H	S	2-N ₃	4-H
	H	H	S	3-I	4-NH ₂
	H	H	O	H	4-NH ₂

5

	H	H	O	H	4-N ₃
	H	H	S	2-F	4-NH ₂
	H	H	S	2-F	4-NO ₂
	H	H	S	2-F	4-N ₃
5	H	H	S	2-Cl	4-NH ₂
	H	H	S	2-Cl	4-NO ₂
	H	H	S	2-Cl	4-N ₃
	4-OMe	H	S	2-Cl	4-NO ₂
	5-OMe	H	S	2-Cl	4-NO ₂
10	6-OMe	H	S	H	4-NO ₂
	6-OMe	H	S	2-Cl	4-NO ₂
	7-OMe	H	S	2-Cl	4-NO ₂
	5-OBenzyl	H	S	2-Cl	4-NO ₂
	6-OBenzyl	H	S	2-Cl	4-NO ₂
15	7-OBenzyl	H	S	2-Cl	4-NO ₂
	5-OMe	6-OMe	S	H	4-NO ₂
	6-OMe	7-OMe	S	H	4-NO ₂
	5-OMe	7-OMe	S	H	4-NO ₂
	4-OMe	H	S	2-Cl	4-NH ₂
20	5-OMe	H	S	2-Cl	4-NH ₂
	6-OMe	H	S	2-Cl	4-NH ₂
	5-OH	H	S	2-Cl	4-NH ₂
	6-OH	H	S	2-Cl	4-NH ₂
	5-OBenzyl	H	S	2-Cl	4-NH ₂
25	H	H	S	6-OSO ₂ CF ₃	3-NH ₂
	H	H	S	3-OSO ₂ CF ₃	4-NH ₂
	6-Me	H	S	3-OSO ₂ CF ₃	4-NH ₂
	H	H	S	5-OSO ₂ CF ₃	2-NH ₂
	7-OBenzyl	H	S	2-Cl	4-NH ₂
30	6-OMe	H	S	H	4-NH ₂
	6-OH	H	S	H	4-NH ₂
	6-Me	H	S	H	4-NH ₂
	5-OMe	7-OMe	S	H	4-NH ₂
	5-OH	7-OH	S	H	4-NH ₂
35	5-OH	7-OMe	S	H	4-NH ₂
	5-OMe	6-OMe	S	H	4-NH ₂

In preferred compounds, as may be apparent from the

above list, R^1 and R^3 will often both be hydrogen but one or other may alternatively be alkoxy or aralkoxy, e.g. methoxy or benzyloxy, whilst at least one of R^2 and R^4 is preferably an amino or a substituted amino group (e.g. -
5 NMe₂) but may alternatively be nitro, azido, hydroxy, halo, or R^2 and R^4 may collectively comprise a combination of such groups substituted at different positions. Other substitution patterns can also be of interest as will be apparent from examples herein disclosed. Although in some
10 cases the nitro and azido substituted compounds themselves exhibit some selective cytotoxic activity, most commonly these derivatives will be used as intermediates for reduction to obtain the corresponding amino compounds. The latter are usually more likely to show a useful level of
15 biological activity.

One especially important compound in the above list is that designated CJM 126 wherein R^1 , R^2 and R^3 are each hydrogen, R^4 is 4-NH₂ and X is sulphur, i.e. 2-(4-amino-
20 phenyl)benzothiazole, including pharmaceutically acceptable salts thereof.

Salts of the compounds of formula I in accordance with the invention may include acid addition salts derived
25 from an acid selected from the group comprising: hydrochloric, hydrobromic, sulphuric, nitric, phosphoric, maleic, salicylic, p-toluenesulphonic, tartaric, citric, lactobionic, formic, malonic, pantothenic, succinic, naphthalene-2-sulphonic, benzenesulphonic, methanesulphonic
30 and ethanesulphonic.

The invention also comprises the use of a 2-arylbenzazole compound as specified above for making a medicament or pharmaceutical composition for use in
35 antitumour therapy, especially in the treatment of breast cancer for example.

As hereinafter described, the invention also includes pharmaceutical compositions or preparations, conveniently

in unit dosage form, for use in antitumour therapy, especially breast cancer therapy, said compositions or preparations comprising as the active substance a 2-arylbenzothiazole or 2-arylbenzoxazole compound as
5 specified above.

In general, salts of compounds such as CJM 126, e.g. hydrochloride, dihydrochloride, methanesulphonic acid and ethanesulphonic acid addition salts, should be equally
10 effective in inhibiting proliferation of breast tumour cells insofar as such salts will dissociate in water or other aqueous media to provide the active antitumour compound. In practice these salts may be the most preferred compounds for making up acceptable pharmaceutical
15 formulations.

Development and Testing

The invention has developed from an initial
20 observation made when carrying out certain routine chemical investigations. During these investigations it was found that the arylbenzothiazole compound 2-(4-aminophenyl)benzothiazole, herein designated CJM 126 and sometimes referred to as ptiazamine, had a very surprising and completely
25 unexpected high and selective activity as an anti-proliferative agent in respect of cultures of MCF-7 mammary carcinoma cells.

This is illustrated in Tables 1 and 2 at the end of
30 the present description which show in vitro test results obtained in various sets of experiments for the cytotoxic activity of this compound CJM 126 against a range of human tumour cell lines, expressed in terms of IC₅₀ values (concentration or dosage required to reduce cell growth or
35 proliferation by 50%) calculated from dose-effect curves plotted for cultures of the cells in question.

It will be seen from Table 1 that the MCF-7 cell line used in the first series of experiments was about 3000

times more sensitive to CJM 126 than other human cell lines tested. This remarkable result has been reconfirmed on several subsequent occasions, as evidenced for example by the results presented in Table 2, using fresh samples and

5 different strains of the MCF-7 cells in later series of experiments. Thus, selective inhibition has been clearly demonstrated, for example against the following human mammary carcinoma cell lines: MCF 7-wt, MDA 468, SKBR3, ZR 75 and MDA 231 with estimated IC_{50} values of about 2.9nM,

10 1.6nM, 25.9nM, 24.7nM, and 17.2nM respectively. Also, a somewhat unusual biphasic dose response relationship was noted. For example, in the first series of experiments referred to above it was found that an initial decrease in cell growth for the MCF-7 cells, which gave the IC_{50} value

15 of approximately 2 ng/ml, was followed at higher doses by a small increase, and then at even greater doses there was a second decrease corresponding to an IC_{50} value of about 2 μ g/ml. This biphasic dose response effect has also been found to be consistently reproducible. Thus, in the later

20 series of experiments 30nM and 100nM CJM 126 resulted in maximum growth inhibition of MCF-7 cells and loss of growth arrest, characterised by rising cell numbers, followed exposure to 1 μ M, 3 μ M and 10 μ M. For MCF 7-ADR cells at low concentrations of CJM 126 growth was negligible under the

25 experimental conditions used (seeding density 2.5×10^2 cells per well). However, at higher concentrations of 3 and 10 μ M, CJM 126 induced proliferation of these MCF cells 7-ADR.

30 Although certain mammary carcinoma cell lines examined did not give a biphasic dose response following CJM 126 exposure, cell lines derived from tumours of other origins appeared generally to be insensitive to the specific inhibitory properties of CJM 126. No effect was

35 observed, for example, on the growth of A431 cells as a result of exposure to CJM 126 over a range of concentrations (0.1nM-10 μ M). Estimated IC_{50} values for some of the other cell lines tested are given in Table 2.

In other in vitro studies it was found that CJM 126 gave an IC_{50} value of 2.7×10^{-5} M against P388 cells, and against P388R8/22 (multidrug resistant) cells it gave an IC_{50} value of 5.2×10^{-5} M. CJM 126 was also tested
5 against the NCI (National Cancer Institute, U.S.A.) panel of human tumour cell lines which does not contain any breast cancer cell lines, and the compound was found to be essentially non-toxic in these tests.

10 In carrying out the above in vitro studies, the method used for conducting the cytotoxicity assays was generally substantially as follows:

Cells were maintained in a continuous logarithmic culture in Dulbecco's medium supplemented with 10%
15 fetal calf serum and penicillin (100 IU/ml) and streptomycin (100 µg/ml). The cells were mildly trypsinized for passage and for use in assays. On day zero, 100µl of trypsinized tumour cells (1×10^4 /ml) were plated in the wells of 96-well flat-
20 bottom microtiter plates. The plates were incubated for 2 days at 37°C and 5% CO₂ in air to allow the cells to adhere and resume exponential growth prior to the addition of drugs.

25 The compounds being tested were dissolved in a small volume of DMSO and diluted to the desired concentration with growth medium so that the final concentration of DMSO did not exceed 0.25%. On day two 50µl of the highest drug concentration was added
30 to the wells of column 12 and from there serially diluted 3-fold to column 1 by serial transfer of 50µl using an 8-channel micropipette. The final volume of column 1 was adjusted to 100µl. No additions were made to the wells of rows A and B, which served as
35 controls. The plates were further incubated for 5 days at 37°C and 5% CO₂ in air. Each compound was tested in duplicate.

On day 7 the test was terminated by the addition of

100µl saline containing 0.002% w/v propidium iodide (Sigma), 0.3% drawing ink (Staedtler "Marromatic 745" - Trade Mark) and 0.5% Triton X-100. The plates were kept at 4°C overnight before reading on an inverted microscope equipped with an automated scanning stage. Fluorescence intensity was measured in arbitrary units by a photomultiplier. An HP-87 computer controlled the movement of the stage and also collected and processed the data from the multiplier.

For each compound tested a dose-response curve was obtained and the IC₅₀ value (the drug concentration at 50% inhibition of cell growth) was calculated.

The remarkably specific effects and activity of CJM 126 noted against human MCF-7 cells in vitro has been found subsequently to extend to in vivo tests and also to a broad range of analogues of CJM 126, as illustrated for example in Tables 3, 4 and 5.

Table 3 presents the results, including estimated IC₅₀ values, of cytotoxicity and proliferation or growth stimulation tests carried out in vitro for different breast cancer cell lines, including MCF7-wt, MDA 468 and MCF 7-ADR, in respect of compound CJM 126 and a range of analogues of CJM 126 identified by reference codes in the first column. The structures of these analogues will be apparent from the subsequent description herein.

The results of some of the in vivo tests carried out, and a comparison of the activity of CJM 126 with that of Mitoxantrone, are set out in Table 4. The results of similar tests carried out to evaluate the activity of CJM 126, and also of a close analogue thereof designated CJM 129, against BO (T61) human mammary carcinomas implanted in NMRI-nu/nu female mice are shown in Table 4 together with results obtained using other antitumour drugs for comparison. The results marked * are all statistically significant (p<0.05). As will be seen, CJM 126 had

significant activity against MCF-7 and BO mammary tumours and there were no signs of severe toxicity following administration of a single i.p. injection of 120 mg/kg CJM 126 (Table 4). Additionally, the relative BO tumour volume was lower following daily treatment of 1mg/kg compared with 10 or 100mg/kg/day (Table 5).

The effect of CJM 126 (0.1, 0.01 and 0.001mg/kg) has also been investigated in nude mice bearing human MDA 468 xenograft tumours. The biphasic response obtained in vitro following exposure of MDA 468 cultures to CJM 126 was also observed in vivo when comparable CJM 126 doses were administered. Preliminary data indicate maximum inhibition to tumour growth following daily treatment of the minimum concentration tested, as can be seen from the summary of results presented in Table 6.

Although high concentrations of CJM 126 were found actually to stimulate cell proliferation in some cases, experiments have demonstrated that a continued exposure to CJM 126 is absolutely essential to maintain this stimulation of growth effect which is observed, for example, in the 2nd phase of the MCF 7-wt response. Removal of CJM 126 led not merely to loss of proliferative potential but also to a decline in cell numbers. In contrast, it was found that cell growth following exposure to nM CJM 126 concentrations (phase 1) remained arrested after drug removal and an observed IC₅₀ value of 2.18nM was not significantly affected.

In these in vitro assays, it was noted that serum factors appear to play a significant role in the 2nd phase of the CJM 126 dose response. Thus, in the presence of 1% FCS, no proliferation of colonies was observed following exposure to 3, 10, 30µM CJM 126, and growth inhibition was maintained. Moreover, enhanced potency was detected with estimated IC₅₀ values in the picomolar range.

In other experiments it was established that growth

of MCF 7-wt cell cultures supplemented with 1% FCS was stimulated 240%, 346% and 397% by 5, 50 and 100ng/ml EGF respectively, but CJM 126 was able to reverse this effect and exquisitely arrest the growth of these cells. However, the dose response profiles and IC_{50} values did not differ significantly from MCF 7-wt cultures supplemented by 1% FCS in the absence of EGF. It was also noted that the presence of EGF was unable to rescue the 2nd phase of the dose response, and inhibition of growth was greater than 85% for all concentrations of CJM 126 between and including 1nM-10 μ M. Similarly, in MCF 7-wt cultures supplemented with 10% FCS, the dose response to CJM 126 was not significantly altered by inclusion of EGF (5, 50 and 100ng/ml) in experimental media.

15

It was also observed that CJM 126 (1 μ M, 10 μ M and 100 μ M) showed no ability to displace iodinated EGF from EGFR. In addition, Western blot analyses utilizing an anti-phosphotyrosine primary antibody demonstrated no apparent inhibition of unstimulated or of EGF stimulated EGFR tyrosine kinase activity, and at present the mode of action of the CJM 126 compound in these biological systems is not known.

25

In the course of studying the in vivo activity it was found that the lethal dose value LD_{50} of CJM 126, i.e. the dose that was lethal to 50% of the animals tested, was about 125mg/kg in male DBA/2 mice when administered as a single i.p. dose. With daily administration over 3 consecutive days the LD_{50} dose was >31mg/kg/day.

30

Preparative Methods

In most cases the arylbenzazole compounds of the present invention can readily be synthesised by various routes from easily available starting materials, and by way of example, several such general synthetic routes, designated Route A, Route B, Route C, and Route D, are illustrated in Figure 1 of the accompanying drawings in relation specifically to

arylbenzothiazole compounds. A reduction scheme for converting a nitro substituent of an arylbenzothiazole compound into an amino substituent is also depicted as Route E. Such nitro compounds are often prepared for use
5 as intermediates in producing the corresponding amino compounds which usually may be expected to possess greater cytotoxic activity.

Some of the arylbenzazole compounds of the present
10 invention are also known compounds per se that are already commercially available.

In the general method for Route A, which is also applicable to the synthesis of corresponding benzoxazole compounds,
15 typically a mixture of the 2-aminothiophenol (0.05 Mol.) (or the 2-aminophenyl) and the appropriate benzoic acid derivative (0.05 Mol.), together with polyphosphoric acid (85g), is heated at 190-230°C for 4 hours, cooled and poured into a mixture of 10% aqueous sodium bicarbonate
20 (1000ml) and ice. The solid product may then be collected, washed with water and recrystallized.

In the general method for Route B, typically a mixture of 2-aminothiophenol (0.05 Mol.), the appropriate benzaldehyde
25 (0.05 Mol.) and dimethylsulphoxide (30 ml) is heated to 180°C for 15 minutes, cooled and diluted with water (200 ml). The precipitate is then collected, washed with water and crystallised.

30 In the general method for Route C, assuming for example that R^2 is a nitro group NO_2 , a solution of the 2-aminothiophenol (0.05 Mol.) in pyridine (50 ml) is added slowly to a mixture of the appropriate nitrobenzoyl chloride (0.05 Mol.) also in pyridine (50 ml) at 25°C. The
35 reaction is exothermic and is cooled in an ice-bath. The mixture may then be diluted with water (200 ml) and the products are collected and washed with water.

In the general method for Route D, in a typical procedure

the appropriate substituted thiobenzanilide (1 Mol. equiv.) is finely powdered and mixed with a little ethanol to form a wet paste. A 30% w/v solution of aqueous sodium hydroxide (8 Mol. equiv.) is added and diluted with water to form a suspension/solution of the thiobenzanilide in 10% w/v aqueous sodium hydroxide. Aliquots of this suspension/solution are then introduced dropwise at one minute intervals into a stirred solution of potassium ferricyanide (4 Mol. equiv.) in water at 80-90°C. The reaction mixture is heated for a further 30 minutes, then cooled. The 2-arylbenzothiazole products are collected, washed with water and crystallised.

Where R^2 of the 2-arylbenzazole compound synthesised by any of the above routes (or by any other route) is a nitro group NO_2 , this may generally be reduced and converted into the corresponding amine as follows (Route E):

A mixture of the 2-(nitrophenyl)benzazole compound in question (0.05 Mol,) and stannous chloride dihydrate (0.25 Mol.) in absolute ethanol (200 ml) is stirred and refluxed under nitrogen for 1 to 4 hours. The ethanol is then removed under reduced pressure and the residue is dissolved in ethyl acetate (4 x 100ml). The combined organic phases are next shaken with excess aqueous sodium hydroxide to liberate the free amine bases and dissolve the tin residues. The separated organic phase is washed with water, dried (magnesium sulphate) and the solvent is evaporated. Finally, the products are then crystallised.

EXAMPLES

The preparation of a number of particular compounds which are considered to be of interest for use as active therapeutic substances to inhibit proliferation of at least certain breast cancer cells and which provide examples of preferred embodiments of the invention, including CJM 126

and also analogues thereof, will now be described in more detail. The compound reference codes used in Table 3 are also quoted where applicable. It should be understood, however, that these specific examples are not intended to be construed in any way as a limitation in the scope of the invention.

Example I

10 2-(4-aminophenyl)benzothiazole (Compound CJM 126)

(a) In one particular example, 2-(4-aminophenyl)benzothiazole (Compound CJM 126) was prepared in 57% yield from 2-aminothiophenol and 4-aminobenzoic acid using synthetic Route A, the final product being crystallised from methanol as pale yellow needle crystals having a melting point 155-157°C (aqueous solubility about 3.8µg/ml).

(b) In another preparative example, the same compound 2-(4-aminophenyl)benzothiazole was prepared in 73% yield, as beige needles (m.p. 155-156°C) crystallised from methanol, by reduction with stannous chloride hydrate of 2-(4-nitrophenyl)benzothiazole. The latter was first obtained (as yellow crystals, m.p. 229-231°C, crystallised from methanol or dimethyl formamide) either via Route C (71% yield) from 2-aminothiophenol and 4-nitrobenzoyl chloride, or via Route D (10% yield) from the corresponding thiobenzanilide and potassium ferricyanide.

30 Example IA

Ethanesulphonic acid salt of 2-(4-aminophenyl)benzothiazole (Compound 93003)

35 To prepare the ethanesulphonic acid salt, 2-(4-aminophenyl)benzothiazole (0.66g) in ethyl acetate (100 ml) at 25°C was treated with ethanesulphonic acid (0.338g). The solid was collected and washed with ethyl acetate (3 x 50 ml) followed by diethyl ether (3 x 50 ml). The

ethanesulphonic acid salt (0.89g., yield 90%) had m.p. 211-213°C.

5 Example IB

Dihydrochloride salt of 2-(4-aminophenyl)benzothiazole

To prepare the dihydrochloride salt, 2-(4-amino-
10 phenyl)benzothiazole (1.5g) was dissolved in ethyl acetate (250 ml) and a stream of dry hydrogen chloride was passed through the solution for 20 minutes. The yellow dihydrochloride salt was collected and washed with ethyl acetate followed by ether. The dihydrochloride (1.61g) had
15 a m.p. 267-269°C.

Example IC

20 2-(4-Aminophenyl)benzothiazole methanesulphonic acid salt

To a solution of 2-(4-aminophenyl)benzothiazole (0.5g, 2.21mmol) in ethyl acetate (65ml) was added dropwise methanesulphonic acid (0.215g, 2.21mmol) at room
25 temperature. The reaction mixture was stirred for 30 minutes. The product was collected and washed with hot ethyl acetate to give a pale yellow powder (0.67g, 94%), m.p. 261-262°C; $\nu_{\max}/\text{cm}^{-1}$ 3422, 2880, 2633, 1598, 1487, 1322, 1220, 1149, 1043, 967, 780, 755, 561; δ_{H} (DMSO- d_6)
30 8.10(1H, d, J7.9, 4-H), 8.00(3H, m, 7,2',6'-H), 7.52(1H, t, J7.3, 5-H), 7.42(1H, t, J7.5, 6-H), 7.06(2H, d, J8.5, 3',5'-H), 5.72(3H, br s, NH_3^+), 2.41(3H, s, CH_3).

In vitro assay of this compound gave IC_{50} values of
35 0.0007 μM against MCF 7-wt cells and 0.0024 μM against MDA 468 cells, indicating a very high level of selective cytotoxic activity.

Example II2-(2-Chloro-4-nitrophenyl)-4-methoxybenzothiazole

5 This compound was prepared from 2-chloro-4-nitro-2'-methoxythiobenzanilide according to the general procedure of synthetic Route D. A yellow solid formed was immediately collected by filtration when 2-chloro-4-nitro-2'-methoxythiobenzanilide was added to the solution of
10 potassium ferricyanide. The crude product was recrystallised twice from methanol to give yellow crystals (46%), m.p. 187-188°C; $\nu_{\max}/\text{cm}^{-1}$ 3432, 3097, 2836, 1570, 1521, 1476, 1385, 1347, 1277, 1262, 1041, 890, 773, 743; $\delta_{\text{H}}(\text{CDCl}_3)$ 8.60(1H, d, J8.8, 6'-H), 8.37(1H, d, J2.3, 3'-H), 8.19(1H, dd, J2.3, 8.7, 5'-H), 7.55(1H, d, J8.1, 7-H),
15 7.43(1H, t, J8.1, 6-H), 6.97(1H, d, J7.9, 5-H).

Example III

20

5-Benzyloxy-2-(2-chloro-4-nitrophenyl)benzothiazole

 This compound was prepared from 2-chloro-4-nitro-3'-benzyloxythiobenzanilide, again according to the general
25 procedure of synthetic Route D. The crude reaction product was separated into three fractions by flash column chromatography on silica using EtOAc-hexane-chloroform (1:6:1) as the eluate. The first fraction (52%) was 5-benzyloxy-2-(2-chloro-4-nitrophenyl)benzothiazole, m.p.
30 156-157°C; $\nu_{\max}/\text{cm}^{-1}$ 3449, 1603, 1521, 1342, 1277, 1181, 1055, 829, 737; $\delta_{\text{H}}(\text{CDCl}_3)$ 8.60(1H, d, J8.8, 6'-H), 8.45(1H, d, J2.2, 3'-H), 8.27(1H, dd, J2.3, 8.8 5'-H), 7.88(1H, d, J8.9, 7-H), 7.72(1H, d, J2.4, 4-H), 7.55-7.38(5H, m, Ph-H), 7.26(1H, dd, J2.4, 8.9, 6-H), 5.23(2H,
35 s, CH_2O).

Example IV5,7-Dimethoxy-2-(4-nitrophenyl)benzothiazole

5 This compound was prepared from 4-nitro-3',5'-dimethoxythiobenzanilide, also according to the general procedure of Route D. Recrystallisation from ethanol-EtOAc gave a yellow powder (60%), m.p. 238-239°C; $\nu_{\max}/\text{cm}^{-1}$ 3442, 2949, 1605, 1579, 1528, 1428, 1352, 1312, 1155, 1126, 853,
10 820, 688; $\delta_{\text{H}}(\text{CDCl}_3)$ 8.37(2H, d, J' , 5'-H), 8.26(2H, d, J 9.0, 2',6'-H), 7.24(1H, d, J 2.0, 4-H), 6.58(1H, d, J 2.0, 6-H), 4.01(3H, s, 7-OCH₃), 3.95(3H, s, 5-OCH₃); m/z 316(M⁺), 270(M-OCH₃ and -CH₃).

15 Example V6-Methoxy-2-(4-nitrophenyl)benzothiazole

 This compound was prepared from 4'-methoxy-4-nitrobenzanilide, again according to the general procedure of Route D. The crude product was purified by flash column chromatography using EtOAc-hexane (1:3) as the eluate to give the title compound (62%), m.p. 216-217°C; $\nu_{\max}/\text{cm}^{-1}$ 3448, 1593, 1518, 1342, 1315, 1265, 1219, 1066, 849; δ_{H}
20 (CDCl₃) 8.34(2H, d, J 9.1, 3',5'-H), 8.21(2H, d, J 9.1, 2',6'-H), 8.00(1H, d, J 9.0, 4-H), 7.38(1H, d, J 2.5, 7-H), 7.15(1H, dd, J 2.6, 9.0, 5-H), 1.58(3H, s, CH₃), m/z 289 (M⁺), 271(M-CH₃).

30 Example VI2-(4-amino-2-chlorophenyl)-4-methoxybenzothiazole (Compound 93005)

35 This amine was obtained in 77% yield by reduction of 2-(2-Chloro-4-nitrophenyl)-4-methoxybenzothiazole according to the general procedure of synthetic Route E. Characterisation of the product was as follows:
mp. 148-150°C; $\nu_{\max}/\text{cm}^{-1}$ 3460, 3316, 3205, 2962, 1626,

1602, 1567, 1474, 1441, 1414, 1335, 1257, 1043, 772, 741;
 $\delta_{\text{H}}(\text{CDCl}_3)$ 8.19 (1H, d, J8.6, 6'-H), 7.49 (1H, d, J8.0, 7-H), 7.31 (1H, t, J8.0, 6-H), 6.91 (d, J7.8, 5-H), 6.76 (1H, d, J2.3, 3'-H), 6.64 (1H, dd, J2.3, 8.6, 5'-H), 4.07-4.04
5 (5H, m, OCH_3 , NH_2).

Example VII

2-(4-Amino-2-chlorophenyl)-5-benzyloxybenzothiazole 10 (Compound DF161)

This amine was obtained by reduction of 5-benzyloxy-2-(2-chloro-4-nitrophenyl)benzothiazole, again according to the general procedure of Route E. The crude product was
15 chromatographed on silica using EtOAc-hexane (1:1) as the eluate to give 2-(4-amino-2-chlorophenyl)-5-benzyloxybenzothiazole (90%), m.p. 119-121°C; $\nu_{\text{max}}/\text{cm}^{-1}$ 3456, 3380, 1629, 1599, 1443, 1260, 1171, 1051, 802; $\delta_{\text{H}}(\text{CDCl}_3)$ 8.12(1H, d, J8.6, 6'-H), 7.79(1H, d, J8.7, 7-H), 7.65(1H, d, J2.4,
20 4-H), 7.54-7.36(5H, m, Ph-H), 7.14(1H, dd, J2.5, 8.8, 6-H), 6.81(1H, J2.3, 3'-H), 6.69(1H, dd, J2.3, 8.6, 5'-H), 5.20(2H, s, CH_2O), 3.77(2H, s, NH_2).

Example VIII

25

2-(4-Aminophenyl)-5,7-dimethoxybenzothiazole (Compound DF1620)

This amine was formed by reduction of 5,7-dimethoxy-2-(4-nitrophenyl)benzothiazole, also according to the
30 general procedure of Route E. The crude product was chromatographed on silica using EtOAc-hexane (1:1) as the eluate to give 2-(4-aminophenyl)-5,7-dimethoxybenzothiazole (89%), m.p. 150-152°C; $\nu_{\text{max}}/\text{cm}^{-1}$ 3417, 3326, 3203, 3000,
35 1599, 1580, 1485, 1415, 1219, 1150, 1124, 1035, 824, 807, 631; $\delta_{\text{H}}(\text{CDCl}_3)$ 7.91(2H, d, J8.6, 2',6'-H), 7.16(1H, d, J2.0, 4-H), 7.75(2H, d, J2.0, 6-H), 4.03(2H, s, NH_2), 3.96(3H, s, 7- OCH_3), 3.91(3H, s, 5- OCH_3).

Example IX2-(4-Aminophenyl)-6-methoxybenzothiazole (Compound 93002)

5 This amine was formed by reduction of 6-methoxy-2-(4-nitrophenyl)benzothiazole, also according to the general procedure of Route E. The crude product was chromatographed on silica using EtOAc-hexane (1:1) as the eluate to give 2-(4-aminophenyl)-6-methoxybenzothiazole
10 (92%), m.p. 191-193°C; $\nu_{\max}/\text{cm}^{-1}$ 3454, 1626, 1605, 1465, 1436, 1264, 1222, 825; δ_{H} (DMSO- d_6) 7.79(1H, d, J8.9, 4-H), 7.70(2H, d, J8.6, 2',6'-H), 7.62(1H, d, J2.5, 7-H), 7.05(1H, dd, J2.6, 8.9, 5-H), 6.66(2H, d, J8.6, 3',5'-H), 5.84(2H, s, NH₂), 3.84(3H, s, OCH₃); m/z 256 (M⁺), 241 (M-
15 OCH₃).

Example X2-(4-Amino-2-Chlorophenyl)benzothiazole (Compound 93004)

20

 This amine was formed by reduction of 2-(2-chloro-4-nitrophenyl)benzothiazole, again according to the general procedure of Route E. Recrystallisation from methanol gave yellow crystals (93%), m.p. 92-94°C; $\nu_{\max}/\text{cm}^{-1}$ 3455, 3299,
25 3194, 1630, 1620, 1432, 1312, 1262, 1050, 847, 757, 729, 699, 626; δ_{H} (CDCl₃) 8.10 (1H, d, J8.6, 6'-H), 8.04 (1H, d, J7.7, 4-H), 7.89 (1H, d, J7.9, 7-H), 7.47 (1H, dt, J1.3, 7.2, 5-H), 7.36 (1H, dt, J1.2, 7.3, 6-H), 6.78 (1H, d, J2.4, 3'-H), 6.66 (1H, dd, J2.4, 8.6, 5'-H), 4.02 (2H, s,
30 NH₂).

Example XI2-(4-Aminophenyl)-5,7-dihydroxybenzothiazole (Compound DF162Eb)
35

 To a stirred solution of 2-(4-aminophenyl)-5,7-dimethoxybenzothiazole (0.24 g, 0.838 mmol), obtained from Example VIII, in dry dichloromethane (15ml) under nitrogen

and at -70°C , was added dropwise boron tribromide (1.0 M solution) in dichloromethane (5.9 ml) over 30 minutes. The mixture was kept stirred at -70°C for a further 1 hour, then allowed to warm slowly to room temperature and stirred
5 overnight. The mixture was then again cooled to -70°C and methanol was added dropwise until no further reaction was observed. It was then poured into 8% (W/V) aqueous sodium hydroxide (50ml). After acidification with 5M hydrochloric acid to pH7, the mixture was extracted with dichloro-
10 methane-methanol (4:1) (3 x 80 ml). The combined extracts were dried (MgSO_4) and evaporated under reduced pressure. The crude product was then separated into two fractions by flash column chromatography on silica using EtOAc-hexane (3:2) as the eluant. The second fraction (62%) was the
15 required 2-(4-aminophenyl)-5,7-dihydroxybenzothiazole, m.p. 294°C (dec); $\nu_{\text{max}}/\text{cm}^{-1}$ 3438, 3342, 3216, 1603, 1482, 1456, 1434, 1295, 1159, 1093, 829; δ_{H} ($\text{DMSO}-d_6$) 10.30(1H, s, 7-OH), 9.43(1H, s, 5-OH), 7.70(2H, d, J8.6, 2',6'-H), 6.74(1H, d, J2.0, 4-H), 6.64(2H, d, J8.6, 3',5'-H),
20 6.34(1H, d, J2.0, 6-H), 5.84(2H, s, NH_2).

Example XII

25 2-(4-Aminophenyl)-6-hydroxybenzothiazole

This compound was obtained by demethylation with boron tribromide from 2-(4-aminophenyl)-6-methoxybenzothiazole (see Example IX) in a manner similar to that
30 described above for Example XI. The crude product was chromatographed on silica using EtOAc-hexane (1:1) as the eluate to give the title compound (89%), m.p. $262-263^{\circ}\text{C}$; $\nu_{\text{max}}/\text{cm}^{-1}$ 3487, 3388, 1620, 1460, 1289, 1238, 1175, 830; δ_{H} ($\text{DMSO}-d_6$) 9.71(1H, s, OH), 7.72, 7.66(3H, m, 4.2',6'-H),
35 7.32(1H, d, J2.3, 7-H), 6.91(1H, dd, J2.4, 8.7, 5-H), 6.65(2H, d, J8.6, 3',5'-H), 5.80(2H, s, NH_2); m/z 242 (M_+).

Example XIII2-(4-Aminophenyl)benzoxazole (Compound DF140)

5 This provides an example of a benzaxazole instead of
a benzothiazole. In accordance with synthetic Route A, a
mixture of 2-aminophenol (1.5g, 0.0136mol) and 4-
aminobenzoic acid (1.885g, 0.0136mol) in polyphosphoric
acid (20g) was heated at about 190°C for 4 hours, then
10 cooled and poured into 10% aqueous sodium bicarbonate (400
ml). The product was collected by filtration, washed with
water and recrystallised from methanol-H₂O (10:1) to give
small pale yellow crystals (1.76g, 62%), m.p. 176-178°C
(lit 185°C); v_{\max}/cm^{-1} 3472, 3322, 3186, 1614, 1500, 1454,
15 1312, 1242, 1170, 1066, 830, 742; δ_{H} (DMSO-d₆) 7.87(2H, d,
J8.6, 2', 6'-H), 7.69-7.64(2H, m, 5, 6-H), 7.36-7.28(2H, m,
4, 7-H), 6.70(2H, d, J8.6, 3', 5'-H), 6.02 (2H, s, NH₂).

Example XIV

20

2-(3-Amino-6-trifluoromethylsulphonyloxyphenyl)benzo-
thiazole

2-(3-Azidophenyl)benzothiazole (1g) was added in
25 small portions (5 x 0.2g) to a mixture of trifluoromethane-
sulphonic acid (4 ml), trifluoroacetic acid (5 ml) and
trifluoromethylacetic anhydride (1 ml) at 0°C. After
evolution of nitrogen ceased, the mixture was stirred at
20°C for 18 hours, basified with aqueous ammonia and the
30 products extracted into ethyl acetate. The organic layer
was washed with water, dried (MgSO₄) and evaporated to give
a gum which was separated on silica with hexane-ethyl
acetate (6:4) as eluent. The product obtained was the
title compound (70% yield). (Found: M⁺ 374. C₁₄H₉F₃N₂O₃S₂
35 requires MW 374).

An in vitro assay of this compound against MCF-7 (wild
type) cells gave an IC₅₀ value of 50nM which again
indicates a useful level of activity.

Example XV2-(4-Amino-3-trifluoromethylsulphonyloxyphenyl)benzo-
thiazole

5

This compound was prepared in a similar way to the compound of Example XIV from 2-(4-azidophenyl)benzothiazole. The yield was 12%, (Found: M^+ 374. $C_{14}H_9F_3N_2O_3S_2$ requires MW 374).

10

Example XVI2-(4-Amino-3-trifluoromethylsulphonyloxyphenyl)-6-methyl-
benzothiazole

15

This compound was prepared in a similar way to the compound of Example XIV from 2-(4-azidophenyl)-6-methylbenzothiazole. (Found: M^+ 388. $C_{15}H_{11}F_3N_2O_3S_2$ requires MW 386).

20

Example XVII2-(2-Amino-5-trifluoromethylsulphonyloxyphenyl)benzo-
thiazole

25

This compound was prepared in a similar way to the compound of Example XIV from 2-(2-azidophenyl)benzothiazole (Found: M^+ 374. $C_{14}H_9F_3N_2O_3S_2$ requires MW 374).

30

35

Other analogue compounds or derivatives of interest that have been prepared using Route A comprise

- 2-(4-Dimethylaminophenyl)benzothiazole
- 2-(4-Diethylaminophenyl)benzothiazole
- 5 2-(2-Aminophenyl)benzothiazole
- 2-(2-Fluorophenyl)benzothiazole
- 2-(4-Aminobenzyl)benzothiazole

and using Route B comprise

- 2-(4-Hydroxyphenyl)benzothiazole
- 10 2-(4-Pyridyl)benzothiazole
- 2-[4-(Pyrrolidin-1-yl)phenyl]benzothiazole (Compound 93006)

and using Route C comprise

- 15 2-(3-Nitrophenyl)benzothiazole
- 2-(2-Chloro-4-nitrophenyl)benzothiazole
- 4,4'-Bis(benzothiazol-2-yl)azobenzene

and using Route D comprise

- 2-(2-Chloro-4-nitrophenyl)-6-methoxybenzothiazole
- 20 2-(2-Chloro-4-nitrophenyl)-7-methoxybenzothiazole

and by reduction of the corresponding nitro compound using Route E comprise

- 2-(3-Aminophenyl)benzothiazole (Compound CJM 129)
- 25 2-(4-Amino-2-chlorophenyl)-5-methoxybenzothiazole
- 2-(4-Amino-2-chlorophenyl)-6-methoxybenzothiazole
- 2-(4-Amino-2-chlorophenyl)-7-methoxybenzothiazole

Further analogues or derivatives of interest that have been prepared include

- 30 2-(4-Azidophenyl)benzothiazole
- 2-[4-(Morpholin-4-yl)phenyl]benzothiazole (Compound 93008)
- 2-[4-(Piperidin-1-yl)phenyl]benzothiazole (93007)
- 1-(Benzothiazol-2-yl)-2-(4-dimethylaminophenyl)-ethene
- 35 4,4'-Bis(benzothiazol-2-yl)hydrazobenzene (Compound 126-126)
- 2,2'-Diamino-5,5'-Di-(benzothiazol-2-yl)biphenyl (Compound DF68D)

Commercially available analogue compounds that have been tested and considered to be of interest include 2-(4-aminophenyl)-6-methylbenzothiazole for which in vitro assays against MCF 7-wt cells gave an IC_{50} value of 0.38 μ M and against MDA 468 cells an IC_{50} value of 0.4 μ M.

It will be appreciated that some of the above listed compounds with the more complex molecular structures, for example DF68D and 126-126, may be expected to break down or to be metabolised after administration to a mammal so as to form a simpler 2-arylbenzazole compound that, in use, is the main biologically active component. Also, many of the compounds with nitrogen-containing substituents, such as azido groups and nitro groups, may similarly be converted, in use, within the body to a corresponding active amino compound.

Therapeutic Use

As already indicated, the compounds of this invention have been found to inhibit tumour cell proliferation and to have significant selective antitumour activity, especially in respect of breast cancers. Antitumour activity is evidenced for example by reduction of tumour cell number in mammals bearing breast cancer tumours and a consequent increase in survival time as compared to a control provided by animals which are untreated. Antitumour activity is further evidenced by measurable reduction in the size of solid tumours following treatment with the compounds of this invention compared to the tumours of untreated control animals.

Accordingly, as previously stated the compounds of the present invention are of particular interest for the treatment of breast cancer tumours, and the invention further provides a method for the treatment of a patient suffering from breast cancer. For this purpose, an effective non-toxic amount of the active 2-arylbenzazole compound, such as the 2-(4-aminophenyl)benzothiazole

compound or an acid addition salt or close analogue thereof as hereinbefore defined, may be suitably administered, orally, parenterally (including subcutaneously, intramuscularly and intravenously), or topically. The
5 administration will generally be carried out repetitively at intervals, for example once or several times a day.

The amount of the active compound which is required in order to be effective as an antitumour agent for
10 treating mammals will of course vary and is ultimately at the discretion of the medical or veterinary practitioner treating the mammal in each particular case. The factors to be considered by such practitioner, e.g. a physician, include the route of administration and pharmaceutical
15 formulation; the mammal's body weight, surface area, age and general condition; and the chemical form of the compound to be administered. However, a suitable effective antitumour dose may be in the range of about 1.0 to about 75 mg/kg bodyweight, preferably in the range of about 5 to
20 40mg/kg with most suitable doses being for example in the range of 10 to 30mg/kg. In daily treatment for example, the total daily dose may be given as a single dose, multiple doses, e.g. two to six times per day, or by intravenous infusion for any selected duration. For
25 example, for a 75kg mammal, the dose range could be about 75 to 500mg per day, and it is expected that a typical dose would commonly be about 100mg per day. If discrete multiple doses are indicated, treatment might typically be 50mg of the arylbenzazole compound as hereinbefore defined,
30 given 4 times per day in the form of a tablet, capsule, liquid (e.g. syrup) or injection. On account of the biphasic dose response characteristics of many of these compounds, however, care should be taken, particularly in the initial stages of treatment, to ensure that dosage
35 amounts are not too high.

While it may be possible for the active compound of this invention to be administered alone as the raw chemical, it is preferable to present the active compound

as a pharmaceutical formulation. Formulations of the present invention, for medical use, comprise the active compound together with one or more pharmaceutically acceptable carriers thereof and, optionally, any other
5 therapeutic ingredients. The carrier(s) must be pharmaceutically acceptable in the sense of being compatible with the other ingredients of the formulation and not deleterious to the recipient thereof.

10 The present invention therefore further provides a pharmaceutical formulation comprising an arylbenzazole compound as hereinbefore specified (possibly in the form of a free base or a pharmaceutically acceptable acid addition
15 salt) together with a pharmaceutically acceptable carrier thereof.

The formulations include those suitable for oral, rectal, topical and parenteral (including subcutaneous, intramuscular and intravenous) administration.
20

The formulations may conveniently be presented in unit dosage form and may be prepared by any of the methods well known in the art of pharmacy. All methods include generally the step of bringing the active compound into
25 association with a carrier which constitutes one or more accessory ingredients. Usually, the formulations are prepared by uniformly and intimately bringing the active compound into association with a liquid carrier or with a finely divided solid carrier or with both and then, if
30 necessary, shaping the product into desired formulations.

Formulations of the present invention suitable for oral administration may be presented as discrete units such as capsules, cachets, tablets or lozenges, each containing
35 a predetermined amount of the active compound; as a powder or granules; or a suspension in an aqueous liquid or non-aqueous liquid such as a syrup, an elixir, an emulsion or a draught. The active compound may also be presented as a bolus, electuary or paste.

A tablet may be made by compression or moulding, optionally with one or more accessory ingredients. Compressed tablets may be prepared by compressing, in a suitable machine, the active compound in a free-flowing
5 form such as a powder or granules, optionally mixed with a binder, lubricant, inert diluent, surface active or dispersing agent. Moulded tablets may be made by moulding, in a suitable machine, a mixture of the powdered active compound with any suitable carrier.

10

A syrup may be made by adding the active compound to a concentrated, aqueous solution of a sugar, for example sucrose, to which may be added any accessory ingredient. Such accessory ingredient(s) may include flavourings, an
15 agent to retard crystallisation of the sugar or an agent to increase the solubility of any other ingredient, such as a polyhydric alcohol for example glycerol or sorbitol.

Formulations for rectal administration may be
20 presented as a suppository with a usual carrier such as cocoa butter.

Formulations suitable for parental administration conveniently comprise a sterile aqueous preparation of the
25 active compound which is preferably isotonic with the blood of the recipient.

In addition to the aforementioned ingredients, formulations of this invention, for example ointments,
30 creams and the like, may include one or more accessory ingredient(s) selected from diluents, buffers, flavouring agents, binders, surface active agents, thickeners, lubricants, preservatives (including antioxidants) and the like.

35

From another aspect, the invention thus also comprises use of a benzazole compound as hereinbefore specified for the manufacture of a medical preparation for the treatment of breast cancer tumours.

Table 1. In vitro cytotoxicity of CJM 126 against human cell lines (1st series of experiments)

Cell Line	IC ₅₀ (ng/ml)
MCF-7 mammary carcinoma	2
A204 rhabdomyosarcoma	5943
T24 bladder carcinoma	16876
WiDr colon carcinoma	6008
IGR37 melanoma	23986
HT29 colon carcinoma	7568
A2780 ovarian carcinoma	8826

Table 2. IC₅₀ values for CJM 126 against human cell lines (2nd series of experiments)

Cell Line	IC ₅₀ μM
MCF 7 mammary (ER+)	0.003
MCF 7 mammary (ER+)	0.008
MCF 7-ADR mammary	stimulation > 1μM
ZR 75 mammary (ER+)	0.028
SKBR3 mammary (ER-)	0.026
MDA 468 mammary (ER-/EGFR+)	0.0016
MDA 231 mammary (EGFR+/erbB3+)	.017
T47D mammary (ER-)	44.1
MCF 7-B (ER+)	62.2
MCF 7-T (ER-)	12.87
A204 rhabdomyosarcoma	26
T24 bladder carcinoma	75
WiDr colon	26
IGR 37 melanoma	106
HT 29 colon	33
A2780 ovarian carcinoma	39
A2780 ovarian carcinoma	43
A2780/CisR	37
HX/62 ovarian carcinoma	120
SKOV-3 ovarian carcinoma	64
41M ovarian carcinoma	36
41M/CisP	40
CH 1 ovarian carcinoma	22
CH 1/CisR	33

Table 3 Cytotoxicity (IC₅₀) and growth stimulation of analogues of CJM 126 in breast cancer cell lines *in vitro*

Compound	MCF-wt (μ M)		MDA 468 (μ M)		MCF7-Adr (μ M)		
	A	B	A	B	A	B	C
CJM126	0.003	-	0.0016	-	-	-	>1
DF140	0.11	-	-	>10	-	-	10, 30
DF161	0.001	55.9	0.0007	2.84	-	84.8	3,10,30
93002	0.38	-	-	7.64	-	-	3,10
93003	0.0046	-	0.0024	-	-	-	3,10
93004	0.142	-	0.003	-	-	-	1,3,10
CJM129	-	>10	NT	NT	Inactive		
DF126-126	-	>10	NT	NT	Inactive		
93005	0.063	-	0.0022	NT	0.0465	-	3,10
93006	0.867	-	NT	NT	-	-	3,10
93007	-	2.77	NT	NT	-	-	1,3,10
93008	0.160	-	NT	NT	-	-	3,10
DF162Eb	-	49.65	-	16.3	-	13.9	-
DF162D	-	19.3	0.0035	-	-	20.9	-
DF68D	-	22.3	-	19.3	Inactive		
A	IC ₅₀ at sub-micromolar concentrations						
B	IC ₅₀ at >micromolar concentrations						
C	Concentrations which stimulate cell growth						
NT	Not tested						

Table 4 *In vivo* activity of CJM 126 against MCF-7 mammary carcinoma

Compound	Route	Dose mg/kg/day	BWC (%)	d9	Relative tumour volume (T/C%)				
					d13	d20	d27	d35	d43
CJM 126	i.p.	120	-13	84	47*	38*	27*	27*	43*
Mitoxantrone	i.v.	10	-10	52	32*	38*	26*	29*	29*

Mice: Bln. NMRI - nu/nu (female)

Tumour: MCF-7 implanted s.c. supplemented with oestradiol

Treatment schedule: qd 6, 13, 20 by indicated route.

BWC: Body weight change relative to saline treated animals

Control: Saline

*Statistically significant (p<0.05)

Table 5 *In vivo* activity of CJM 126 and CJM 129 against BO mammary carcinoma

Compound	Route	Dose (mg/kg/day)	BWC	Relative tumour volume (T/C%)				
				d33	d41	d47	d55	d61
CJM 126	i.p.	100	-7	137	77	87	68	52
CJM 126	i.p.	10	-4	107	74	55	70	52
CJM 126	i.p.	1	-5	173	72	25*	67	41*
CJM 129	i.p.	200	-6	78*	45*	32*	33*	36*
CJM 129	i.p.	20	-3	88	77	54	73	48*
CJM 129	i.p.	2	2	127	91	94	95	78
Vincristine	i.p.	1	-6	142	62	61	61	57
Cyclophosphamide	i.p.	150	-4	117	27*	29*	19*	5*
Mitoxantrone	i.v.	10	-5	58*	26*	6*	7*	4*

Treatment schedule : for CJM 126 and CJM 129 qd 27, 34, 41; for vincristine, cyclophosphamide and mitoxantrone qd 27.

CJM 129 is 2-(3-aminophenyl)benzothiazole

For other details: see footnotes to Table 4


SUBSTITUTE SHEET (RULE 26)

Table 6. Activity of CIM 126 against human MDA 468 tumour *in vivo*.

<u>Day</u>	<u>Cross section tumour area (mm)</u>			
	<u>Group</u>			
	<u>Control</u>	<u>.001mg/kg</u>	<u>.01mg/kg</u>	<u>0.1mg/kg</u>
5	95.28 + 4.61	85.54 + 19.44	120.78 + 5.83	97.75 + 9.55
12	62.88 + 8.10	52.93 + 2.65	71.65 + 11.67	71.07 + 7.2
15	86.02 + 8.65	67.81 + 12.92	78.21 + 6.69	82.17 + 10.58
19	118.35 + 37.1	64.8 + 17.35	98.03 + 33.51	88.84 + 3.69
22	119.67 + 19.05	75.12 + 20.05	117.02 + 42.1	105.66 + 5.49
26	130.97 + 20.61	81.53 + 26.16	131.58 + 23.41	124.55 + 7.90
29	170.99 + 51.67	114.65 + 34.57	164.90 + 37.18	182.15 + 14.79
32	191.0 + 48.36	129.06 + 32.08	191.68 + 45.65	196.66 + 18.65
<u>Tumour weight (g)</u>				
	.31 + .09	.19 + .08	.29 + .15	.30 + .07


CLAIMS

1. For use in therapy, a compound that is a benzazole compound represented by the structural formula I below, or
5 a pharmaceutically acceptable salt thereof,



characterised in that

- 15 X is S or O;
R¹ and R³ are each independently hydrogen, alkyl, hydroxyl, alkoxy or aralkoxy
Z is aryl;
- 20 subject to the proviso that alkyl groups when present as such in the compound or as a moiety in other groups such as alkoxy are each composed of less than 6 carbon atoms.
- 25 2. A 2-arylbenzazole compound as claimed in Claim 1 for use as a therapeutic agent wherein Z is



- 35 where n = 0 or 1,
R² is hydrogen, NH₂, NO₂, N₃, halogen or an alkyl or substituted alkyl oxysulphonyl group;

R^4 is hydrogen, NO_2 , N_3 , pyrrolidino, piperidino, morpholino or NR^5R^6 where

5 R^5 and R^6 each represent hydrogen or alkyl;

with the further proviso that R^2 and R^4 are not both hydrogen.

10 3. A compound as claimed in Claim 2 for use as an active therapeutic substance further characterised in that at least one of R^2 and R^4 is an amino or a substituted amino group. .

15 4. A 2-arylbenzazole compound as claimed in Claim 2 for use as a therapeutic agent wherein Z is



5. A compound as claimed in Claim 4 for use as an active therapeutic substance wherein R^2 is 2-Cl.

25

6. A compound as claimed in any of Claims 2 to 5 for use as an active therapeutic substance wherein R^1 is hydrogen and R^3 is alkoxy or benzyloxy.

30 7. A compound as claimed in any of Claims 2 to 6 further characterised by at least one of the following features:

- (a) at least some alkyl groups when present as such or as a moiety in other groups such as alkoxy are methyl or ethyl;
- 35 (b) halo substituents, when present, are selected from fluorine and chlorine;
- (c) R^2 is hydrogen or 2-halogen, and R^4 is amino.

8. A 2-arylbenzazole compound as claimed in any of the

preceding claims for use as an active therapeutic substance, said compound being a substituted 2-phenylbenzothiazole compound.

- 5 9. A compound as claimed in Claim 2 for use as an active therapeutic substance, said compound being a substituted 2-phenylbenzazole compound characterised in that the combination of substituents R^1 , R^2 , R^3 , R^4 and X is selected from the following combinations:

10

	<u>R^1</u>	<u>R^3</u>	<u>X</u>	<u>R^2</u>	<u>R^4</u>
	H	H	S	H	4-NH ₂
	H	H	S	3-NH ₂	4-H
	H	H	S	2-NH ₂	4-H
15	H	H	S	H	4-NMe ₂
	H	H	S	H	4-NEt ₂
	H	H	S	H	4-Pyrrolidino
	H	H	S	H	4-Piperidino
	H	H	S	H	4-Morpholino
20	H	H	S	H	4-NO ₂
	H	H	S	3-NO ₂	4-H
	H	H	S	2-NO ₂	4-H
	H	H	S	H	4-N ₃
	H	H	S	3-N ₃	4-H
25	H	H	S	2-N ₃	4-H
	H	H	S	3-I	4-NH ₂
	H	H	O	H	4-NH ₂
	H	H	O	H	4-N ₃
	H	H	S	2-F	4-NH ₂
30	H	H	S	2-F	4-NO ₂
	H	H	S	2-F	4-N ₃
	H	H	S	2-Cl	4-NH ₂
	H	H	S	2-Cl	4-NO ₂
	H	H	S	2-Cl	4-N ₃
35	4-OMe	H	S	2-Cl	4-NO ₂
	5-OMe	H	S	2-Cl	4-NO ₂
	6-OMe	H	S	H	4-NO ₂
	6-OMe	H	S	2-Cl	4-NO ₂

	7-OMe	H	S	2-Cl	4-NO ₂
	5-OBenzyl	H	S	2-Cl	4-NO ₂
	6-OBenzyl	H	S	2-Cl	4-NO ₂
	7-OBenzyl	H	S	2-Cl	4-NO ₂
5	5-OMe	6-OMe	S	H	4-NO ₂
	6-OMe	7-OMe	S	H	4-NO ₂
	5-OMe	7-OMe	S	H	4-NO ₂
	4-OMe	H	S	2-Cl	4-NH ₂
	5-OMe	H	S	2-Cl	4-NH ₂
10	6-OMe	H	S	2-Cl	4-NH ₂
	5-OH	H	S	2-Cl	4-NH ₂
	6-OH	H	S	2-Cl	4-NH ₂
	5-OBenzyl	H	S	2-Cl	4-NH ₂
	H	H	S	6-OSO ₂ CF ₃	3-NH ₂
15	H	H	S	3-OSO ₂ CF ₃	4-NH ₂
	6-Me	H	S	3-OSO ₂ CF ₃	4-NH ₂
	H	H	S	5-OSO ₂ CF ₃	2-NH ₂
	7-OBenzyl	H	S	2-Cl	4-NH ₂
	6-OMe	H	S	H	4-NH ₂
20	6-OH	H	S	H	4-NH ₂
	6-Me	H	S	H	4-NH ₂
	5-OMe	7-OMe	S	H	4-NH ₂
	5-OH	7-OH	S	H	4-NH ₂
	5-OH	7-OMe	S	H	4-NH ₂
25	5-OMe	6-OMe	S	H	4-NH ₂

10. 2-(4-aminophenyl)benzothiazole for use as an active therapeutic substance or a pharmaceutically acceptable salt thereof.

30

11. A compound as claimed in Claim 1 for use as an active therapeutic substance, said compound being one of the following:

- 35
- (1) 2-(4-Dimethylaminophenyl)benzothiazole
 - (2) 2-(4-Diethylaminophenyl)benzothiazole
 - (3) 2-(2-Aminophenyl)benzothiazole
 - (4) 2-(2-Fluorophenyl)benzothiazole
 - (5) 2-(4-Aminobenzyl)benzothiazole
 - (6) 2-(4-Hydroxyphenyl)benzothiazole

- (7) 2-(4-Pyridyl)benzothiazole
- (8) 2-[4-(Pyrrolidin-1-yl)phenyl]benzothiazole
- (9) 2-(3-Nitrophenyl)benzothiazole
- (10) 2-(2-Chloro-4-nitrophenyl)benzothiazole
- 5 (11) 6-Methoxy-2-(4-nitrophenyl)benzothiazole
- (12) 2-(2-Chloro-4-nitrophenyl)-6-methoxybenzothiazole
- (13) 2-(2-Chloro-4-nitrophenyl)-7-methoxybenzothiazole
- (14) 2-(2-Chloro-4-nitrophenyl)-4-methoxybenzothiazole
- (15) 2-(3-Aminophenyl)benzothiazole
- 10 (16) 2-(4-Amino-2-chlorophenyl)benzothiazole
- (17) 2-(4-Amino-2-chlorophenyl)-4-methoxybenzothiazole
- (18) 2-(4-Amino-2-chlorophenyl)-5-methoxybenzothiazole
- (19) 2-(4-Amino-2-chlorophenyl)-6-methoxybenzothiazole
- (20) 2-(4-Amino-2-chlorophenyl)-7-methoxybenzothiazole
- 15 (21) 2-(4-Azidophenyl)benzothiazole
- (22) 2-[4-(Morpholin-4-yl)phenyl]benzothiazole
- (23) 2-[4-(Piperidin-1-yl)phenyl]benzothiazole
- (24) 4,4'-Bis(benzothiazol-2-yl)azobenzene
- (25) 1-(Benzothiazol-2-yl)-2-(4-dimethylaminophenyl)-
20 ethene
- (26) 4,4'-Bis(Benzothiazol-2-yl)hydrazobenzene
- (27) 2,2'-Diamino-5,5'-Di-(benzothiazol-2-yl)biphenyl
- (28) Ethanesulphonic acid salt of 2-(4-aminophenyl)-
benzothiazole
- 25 (29) Dihydrochloride salt of 2-(4-aminophenyl)benzo-
thiazole
- (30) 2-(4-Aminophenyl)benzothiazole methanesulphonic
acid salt

30 12. A compound as claimed in any of the preceding claims
for use as an active therapeutic substance characterised in
that it is an acid addition salt derived from an acid
selected from the group comprising: hydrochloric,
hydrobromic, sulphuric, nitric, phosphoric, maleic,
35 salicylic, p-toluenesulphonic, tartaric, citric,
lactobionic, formic, malonic, pantothenic, succinic,
naphthalene-2-sulphonic, benzenesulphonic, methanesulphonic
and ethanesulphonic.

13. A process for preparation of a compound having the general structure defined in Claim 2 wherein R^2 and/or R^4 is amino, said process being characterised by the fact that it comprises the steps of preparing the corresponding nitro substituted compound, i.e. the compound with the same structural formula except that R^2 and/or R^4 is a nitro group, and then reducing said nitro group(s) to amino.

14. A compound as claimed in any one of Claims 1 to 12 for therapeutic use in treating breast cancer in mammals.

15. A pharmaceutical formulation for medical use comprising, as the active compound, a compound as claimed in any one of Claims 1 to 12 together with a pharmaceutically acceptable carrier therefor.

16. A medical preparation containing a therapeutically effective non-toxic amount of a compound as claimed in any one of Claims 1 to 12 and a pharmaceutically inert excipient.

17. A pharmaceutical preparation in unit dosage unit form for administration to obtain a therapeutic effect as an antitumour agent in treating mammals, said preparation comprising, per dosage unit, a therapeutically-effective non-toxic amount of a compound as set forth in any one of Claims 1 to 12.

18. Use of a compound as claimed in any one of Claims 1 to 12 for the manufacture of a medical preparation for the treatment of tumours in mammals.

19. Use as claimed in Claim 18 wherein the medical preparation is for inhibiting the growth or proliferation of human breast cancer cells.

20. A method of treating a mammal suffering from cancer so as to inhibit or reduce cancer cell growth, especially growth or proliferation of breast cancer cells, said method

comprising administering to said mammal an effective antitumour composition wherein the active component is a benzazole compound as claimed in any one of Claims 1 to 12.

5

10

15

20


25

30


35

FIG. 1


Route A




Route B




Route C



Route D



Route E



SUBSTITUTE SHEET (RULE 26)

INTERNATIONAL SEARCH REPORT

International application No.

GB 94/01883

A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 A61K31/425 C07D277/66 C07D263/56 A61K31/42

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 A61K C07D

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	CHEMICAL ABSTRACTS, vol. 71, no. 5, 4 August 1969, Columbus, Ohio, US; abstract no. 22057s, W LEE ET AL 'Induction of increased benzopyrene hydroxylase activity by 2-phenylbenzothiazoles and related compounds' page 323 ; see abstract & CANCER RES., vol.28, no.12, 1968 pages 2539 - 2544 --- -/--	1-20

☒ Further documents are listed in the continuation of box C.☒ Patent family members are listed in annex.

* Special categories of cited documents :

- * "A" document defining the general state of the art which is not considered to be of particular relevance
- * "E" earlier document but published on or after the international filing date
- * "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- * "O" document referring to an oral disclosure, use, exhibition or other means
- * "P" document published prior to the international filing date but later than the priority date claimed

* "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

* "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

* "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

* "&" document member of the same patent family

Date of the actual completion of the international search

11 November 1994

Date of mailing of the international search report

21. 11. 94

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax (+31-70) 340-3016

Authorized officer

Henry, J

INTERNATIONAL SEARCH REPORT

International application No.
/GB 94/01883

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	CHEMICAL ABSTRACTS, vol. 113, no. 24, 10 December 1990, Columbus, Ohio, US; abstract no. 221786q, HU SHENGZHI ET AL 'Crystal structures and antitumor activity of 2-(2'pyridyl)benzothiazole and its organotin complex' page 661 ; see abstract & INORG. CHIM. ACTA, vol.173, no.1, 1990 pages 1 - 4 ---	1,14-20
X	PATENT ABSTRACTS OF JAPAN vol. 10, no. 208 (C-361) 22 July 1986 & JP,A,61 105 975 (TAKEDA CHEM. IND. LTD) 13 March 1986 see abstract ---	1-3,8, 14-20
X	PATENT ABSTRACTS OF JAPAN vol. 13, no. 402 (C-633) 6 September 1989 & JP,A,01 146 875 (KANEBO LTD) 8 June 1989 see abstract ---	1,2,8, 15,16
X	JOURNAL OF MEDICINAL CHEMISTRY, vol.13, no.5, September 1970, WASHINGTON US pages 1012 - 1013 STIG AKERFELDT 'Studies on the in vivo antiviral effects of benzothiazole derivatives against various influenza a2 strains' see the whole document ---	1-4,15, 16
X	PATENT ABSTRACTS OF JAPAN vol. 6, no. 90 (C-104) 27 May 1982 & JP,A,57 021 375 (YAMANOUCHI PHARMCEUT.CO LTD) 4 February 1982 see abstract ---	1,2,8, 15,16
X	JOURNAL OF MEDICINAL CHEMISTRY, vol.13, no.4, July 1970, WASHINGTON US pages 697 - 704 THEODORE H. HASKELL ET AL 'Neuramidinase inhibition and viral chemotherapy' see the whole document ---	1,2,15, 16
X	US,A,2 780 628 (HERSCHEL D. PORTER) 5 February 1957 se column 6, lines 54,55 see the whole document ---	1-4, 9-11,15, 16

-/--

INTERNATIONAL SEARCH REPORT

International application No.

GB 94/01883

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	GB,A,2 164 648 (CIRD) 26 March 1986 see claims ----	1,2,15, 16
X	US,A,4 861 897 (JEFFERY B. PRESS ET AL) 29 August 1989 see column 2, line 39 - column 2, line 50; claims ----	1,2,15, 16
X	GB,A,1 093 355 (TWYFORD LABORATORIES LIMITED) 29 November 1967 see the whole document ----	1,2,15, 16
X	GB,A,1 080 246 (TWYFORD LABORATORIES LIMITED) 23 August 1967 see the whole document ----	1,2,15, 16
X	FR,A,1 255 312 (THE CROOKES LABORATORIES LIMITED) 30 January 1961 see the whole document ----	1,2,15, 16
X	FR,A,2 684 377 (SYNTHELABO) 4 June 1993 see claims -----	1,2,15, 16

INTERNATIONAL SEARCH REPORT

International application No.

PCT/GB94/01883

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
Although claim 20 is directed to a method of treatment of the human body,
the search has been carried out and based on the alleged effects of the
compounds
2. ☐ Claims Nos.:
because they relate to parts of the international application that do not comply with the prescribed requirements to such
an extent that no meaningful international search can be carried out, specifically:
The wording Z=aryl is too broadly formulated to permit an adequate search. For
this reason the search has essentially been restricted to compounds of formula
I supported by the examples.
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all
searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment
of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report
covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is
restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

information on patent family members

Patent Application No

PCT/GB 94/01883

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
US-A-2780628		NONE	
GB-A-2164648	26-03-86	LU-A- 85544	03-04-86
		AT-B- 395714	25-02-93
		AU-B- 578310	20-10-88
		AU-A- 4755985	27-03-86
		AU-B- 593838	22-02-90
		AU-A- 6299586	08-01-87
		BE-A- 903254	18-03-86
		BE-A- 904421	30-06-86
		CA-A- 1256862	04-07-89
		CA-A- 1256804	04-07-89
		CH-A- 665841	15-06-88
		CH-A- 675241	14-09-90
		DE-A- 3533308	27-03-86
		FR-A,B 2570377	21-03-86
		GB-A,B 2197320	18-05-88
		JP-A- 61085360	30-04-86
		NL-A- 8502557	16-04-86
		SE-B- 461466	19-02-90
		SE-A- 8504291	20-03-86
		SE-A- 8800949	16-03-88
		US-A- 4920140	24-04-90
		US-A- 5059621	22-10-91
		US-A- 5260295	09-11-93
		US-A- 4740519	26-04-88
		US-A- 5288744	22-02-94
US-A-4861897	29-08-89	NONE	
GB-A-1093355		NONE	
GB-A-1080246		NONE	
FR-A-1255312		NONE	
FR-A-2684377	04-06-93	NONE	